

121 Everett Road
Albany, NY 12205
518-453-9088

KYLE R FLIK, MD
kyleflik@yahoo.com

POSTERIOR SHOULDER STABILIZATION

Weeks 0-3:

- Abduction Sling
- Isometrics in sling
- Grip strengthening

Weeks 3-6:

- Restrict to FF 90°/IR to stomach PROM → AAROM → AROM
- ER with arm at side as tolerated
- Begin isometrics with arm at side – FF/ER/IR/ABD/ADD
- Start scapular motion exercises (traps/rhomboids/lev. scap/etc)
- No cross-arm adduction, follow ROM restrictions
- Heat before treatment, ice after treatment per therapist's discretion

Weeks 6-12:

- Increase ROM to within 20° of opposite side; no manipulations per therapist; encourage patients to work on ROM on a daily basis
- Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilizers
- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- Closed chain exercises

Months 3-12:

- Advance to full ROM as tolerated
- Begin eccentrically resisted motions, plyometrics (ex. Weighted ball toss), proprioception (es. body blade)
- Begin sports related rehab at 3 months, including advanced conditioning
- Return to throwing at 4 ½ months
- Throw from pitcher's mound at 6 months
- MMI is usually at 12 months post op