

orthoNY

www.orthony.com
121 Everett Road
Albany, NY 12205
518-453-9088

Adam G. Suslak, MD
adam.suslak@gmail.com

Post-Operative Rehabilitation Guidelines for Anterior Cruciate Ligament Reconstruction

- 0-2 Weeks: WBAT, Brace locked at 0 degrees for ambulation and sleeping
ROM: 0-90° with emphasis on full extension
Patella mobilization
SLR supine with brace locked at 0 degrees, Quad Sets
Ankle Pumps
Short crank (90mm) ergometry
- 2-6 Weeks: Brace: unlocked @ 2 weeks (if quad control adequate)
discontinued @ 3 weeks if quad adequate for allograft,
six week for autograft
or six weeks for ACL with meniscus repair
No weight bearing past 90° for ACL with meniscus repair
D/C crutches when gait is non-antalgic (six weeks with meniscus repair)
ROM: 0-125 degrees (Maintain full extension)
Active knee extension to 40 degrees
Standard (170mm) ergometry (if knee ROM > 115 degrees)
Leg Press (80-0 degree arc)
Mini Squats / Weight Shifts
Proprioception training
Initiate Step Up program
- 6-14 Weeks: Progressive Squat program
Initiate Step Down program
Leg Press, Lunges
Isotonic Knee Extensions (90-40 degrees, closed chain preferred)
Agility exercises (sport cord)
Versaclimber/Nordic Track
Retrograde treadmill ambulation
- 14-22 weeks: Begin forward running (treadmill) program when 8" step down satisfactory
Continue Strengthening & Flexibility program
Advance Sports-Specific Agility Drills
Start Plyometric program
- > 22 weeks: Advance Plyometric program, Return to Sport (MD Directed)