

Evan D. Boyd, MD
Knee, Shoulder, & Sports Medicine Surgeon
518-489-2663 ext 5215

REHABILITATION PROTOCOL: Pectoralis Major Repair

This protocol is intended to guide clinicians and patients through the post-operative course for a pectoralis major repair. This protocol is time based (dependent on tissue healing) as well as criterion based. Specific intervention should be based on the needs of the individual and should consider exam findings and clinical decision making. If you have questions, contact the referring physician.

Considerations for the Post-Operative Pectoralis Major Repair Rehabilitation Program

One major factor that influences the post-operative pectoralis major repair rehabilitation outcome is type of repair. It is recommended that clinicians collaborate closely with the referring physician to establish if the repair is bone-tendon, tendon-tendon or muscle-tendon which will dictate soft tissue time constraints.

Post-operative Complications:

If the patient develops a fever, unresolving numbness/tingling, excessive drainage from the incision, uncontrolled pain or any other symptoms you have concerns about, you should contact the referring physician. Special care should be taken to monitor an incision in the axillary area due to increased risk of bacterial and moisture buildup

Evan D. Boyd, MD
 Knee, Shoulder, & Sports Medicine Surgeon
 518-489-2663 ext 5215

PHASE I: IMMEDIATE POST-OP (WEEKS 0-3 WEEKS AFTER SURGERY)

Rehab Goals	<ul style="list-style-type: none"> ▪ Protect surgical repair ▪ Reduce swelling, minimize pain ▪ Promote scar mobility ▪ Maintain UE ROM in elbow, hand and wrist ▪ Gradually increase shoulder PROM ▪ Minimize muscle inhibition ▪ Patient education
Sling	<ul style="list-style-type: none"> ▪ Neutral or internal rotation ▪ Use at night while sleeping
Interventions	<p>Swelling Management</p> <ul style="list-style-type: none"> ▪ Ice, compression <p>Range of motion/Mobility (Week 2)</p> <ul style="list-style-type: none"> ▪ PROM: <ul style="list-style-type: none"> ○ ER: Neutral in 0 degrees ADD progressing 5 degrees each week ○ Flex: 45 degrees progressing 5-10 degrees each week ○ ABD: 30 degrees progressing 5 degrees each week ▪ AROM: Elbow, wrist and hand <p>Soft Tissue Mobilization</p> <ul style="list-style-type: none"> ▪ Scar massage (once scar is closed and dry) <p>Strengthening (Week 3)</p> <ul style="list-style-type: none"> ▪ Periscapular: inferior glide (< 35 degrees ABD), low row ▪ Ball squeeze
Criteria to Progress	<ul style="list-style-type: none"> ▪ PROM ER @ 0 degrees ADD 5 degrees ▪ PROM Flex 50 degrees ▪ PROM ABD 35 degrees ▪ Palpable muscle contraction felt in scapular and shoulder musculature ▪ No complications with Phase I

PHASE II: INTERMEDIATE POST-OP (4-6 WEEKS AFTER SURGERY)

Rehab Goals	<ul style="list-style-type: none"> ▪ Protect surgical repair ▪ Reduce swelling, minimize pain ▪ Gradually increase shoulder PROM ▪ Initiate shoulder AAROM ▪ Minimize muscle inhibition ▪ Improve scapular muscle activation ▪ Patient education
Sling	<ul style="list-style-type: none"> ▪ Neutral or internal rotation

Evan D. Boyd, MD
 Knee, Shoulder, & Sports Medicine Surgeon
 518-489-2663 ext 5215

	<ul style="list-style-type: none"> ○ Bone-tendon repairs can begin weaning out of the sling at 4 weeks ○ Tendon-tendon or muscle-tendon repairs should begin to wean at 5-6 weeks
Precautions	<ul style="list-style-type: none"> ▪ No lifting of objects ▪ No supporting of body weight with hands
Interventions *Continue with Phase I interventions	<p>Swelling Management</p> <ul style="list-style-type: none"> ▪ Ice, compression <p>Range of motion/Mobility</p> <ul style="list-style-type: none"> ▪ PROM: <ul style="list-style-type: none"> ○ ER: increase 5 degrees each week ○ Flex: continue to increase 5-10 degrees each week ○ ABD: continue to increase 5 degrees each week ▪ AAROM: Active assistive shoulder flexion, shoulder flexion with cane, cane external rotation stretch, washcloth press <p>Strengthening (Week 3)</p> <ul style="list-style-type: none"> ▪ Shoulder: Submaximal isometrics: ABD, ext, ER (no IR) ▪ Periscapular: scap retraction, prone scapular retraction, standing scapular setting, supported scapular setting
Criteria to Progress	<ul style="list-style-type: none"> ▪ PROM ER @ 0 degrees ADD 20 degrees ▪ PROM Flex 65-85 degrees ▪ PROM ABD 50 degrees ▪ Minimal substitution patterns with AAROM ▪ No complications with Phase II

PHASE III: INTERMEDIATE POSTOP CONTINUED (7-8 WEEKS AFTER SURGERY)

Rehab Goals	<ul style="list-style-type: none"> ▪ Gradually increase shoulder PROM/AAROM ▪ Initiate shoulder AROM ▪ Promote scar mobility ▪ Gradually increase muscle strength ▪ Patient education
Sling	<ul style="list-style-type: none"> ▪ Discontinue
Weightbearing	<ul style="list-style-type: none"> ▪ Can begin weightbearing through surgical side
Additional Interventions *Continue with Phase I-II Interventions	<p>Range of Motion/Mobility</p> <ul style="list-style-type: none"> ▪ PROM <ul style="list-style-type: none"> ○ ER: increase 5 degrees each week to full ○ Flex: continue to increase 5-10 degrees each week to full ○ ABD: continue to increase 5 degrees each week to full ▪ AAROM: Seated shoulder elevation with cane, seated incline table slides, ball roll on wall, wall climbs ▪ AROM: Supine flexion, salutes, supine punch

Evan D. Boyd, MD
 Knee, Shoulder, & Sports Medicine Surgeon
 518-489-2663 ext 5215

	<p>Strengthening</p> <ul style="list-style-type: none"> ▪ Shoulder: Submaximal flex isometrics, side-lying external rotation ▪ Periscapular: Row on physioball, shoulder extension on physioball, resistance band shoulder extension, resistance band seated rows, rowing, lawn mowers, robbery, serratus punches ▪ Elbow: Biceps curl, resistance band bicep curls and triceps <p>Soft Tissue Mobilization</p> <ul style="list-style-type: none"> ▪ Scar mobilization <p>Motor Control</p> <ul style="list-style-type: none"> ▪ External rotation in scaption and Flex 90 degrees (rhythmic stabilization) <p>Stretching</p> <ul style="list-style-type: none"> ▪ Sidelying horizontal ADD, sleeper stretch
Criteria to Progress	<ul style="list-style-type: none"> ▪ PROM ER @ 0 degrees ADD 30 degrees ▪ PROM Flex 75-105 degrees ▪ PROM ABD 60 degrees ▪ Minimal substitution patterns with AROM ▪ No complications with Phase III

PHASE IV: TRANSITIONAL POSTOP (9-14 WEEKS AFTER SURGERY)

Rehab Goals	<ul style="list-style-type: none"> ▪ Restore full shoulder PROM/AROM (week 12-14 for bone-tendon) ▪ Gradually increase muscle strength ▪ Initiate shoulder IR/pec major isometrics ▪ Patient education
Additional Interventions *Continue with Phase I-III interventions	<p>Range of Motion/Mobility</p> <ul style="list-style-type: none"> ▪ PROM: Full ▪ AROM: Full <p>Strengthening</p> <ul style="list-style-type: none"> ▪ Shoulder: Submaximal IR isometrics, submaximal pectoralis isometrics (starting in a shortened position; progressing towards a more lengthened position), standing external rotation w/ resistance band, external rotation, side lying ABD→standing ABD ▪ Periscapular: Push-up plus on knees, prone shoulder extension Is, tripod, pointer <p>Motor Control</p> <ul style="list-style-type: none"> ▪ PNF – D1 diagonal lifts (concentric to begin, then eccentric; manual resistance progressing to resistance bands)

Evan D. Boyd, MD
 Knee, Shoulder, & Sports Medicine Surgeon
 518-489-2663 ext 5215

	<ul style="list-style-type: none"> ▪ PNF – D2 diagonal lifts (concentric to begin, then eccentric; manual resistance progressing to resistance bands) ▪ Quadruped alternating isometrics ▪ Ball stabilization on wall <p>Stretching</p> <ul style="list-style-type: none"> ▪ Triceps and lats
Criteria to Progress	<ul style="list-style-type: none"> ▪ Full pain-free PROM/AROM ▪ Minimal to no substitution patterns with shoulder AROM ▪ Performs all exercises demonstrating symmetric scapular mechanics

PHASE V: ADVANCED STRENGTHENING (14-20 WEEKS AFTER SURGERY)

Rehab Goals	<ul style="list-style-type: none"> ▪ Restore full shoulder PROM/AROM (week 14-16 for tendon-tendon or muscle-tendon) ▪ Gradually increase muscle strength through the full ROM ▪ Do not overstress healing tissue ▪ Patient education
Additional Interventions *Continue with Phase II-IV interventions	<p>Range of motion/mobility</p> <ul style="list-style-type: none"> ▪ PROM: Full ▪ AROM: Full <p>Stretching</p> <ul style="list-style-type: none"> ▪ Shoulder: Standing internal rotation w/ resistance band, internal rotation, pectoralis isotonic, counter push-ups→push-ups, lat pull downs ▪ Periscapular: Resistance band forward punch, forward punch, T and Y, “T” exercise, “W” exercise, resistance band W’s, dynamic hug, resistance band dynamic hug <p>Motor Control</p> <ul style="list-style-type: none"> ▪ Field goals, wall slides w/ resistance band <p>Stretching</p> <ul style="list-style-type: none"> ▪ Hands behind head, IR behind back with towel, doorway series (gentle stretch only)
Criteria to Progress	<ul style="list-style-type: none"> ▪ Full pain-free PROM and AROM ▪ Minimal to no substitution patterns with shoulder AROM ▪ Performs all exercises demonstrating symmetric scapular mechanics ▪ Pain < 2/10

PHASE VI: EARLY RETURN TO SPORT (5-6 MONTHS AFTER SURGERY)

Evan D. Boyd, MD
 Knee, Shoulder, & Sports Medicine Surgeon
 518-489-2663 ext 5215

Rehab Goals	<ul style="list-style-type: none"> ▪ Maintain pain-free ROM ▪ Continue strengthening and motor control exercises ▪ Enhance functional use of upper extremity
Additional Interventions *Continue with Phase II-V interventions	<p>Strengthening</p> <ul style="list-style-type: none"> ▪ Shoulder: External rotation at 90 degrees, internal rotation at 90 degrees, resistance band standing external rotation at 90 degrees, resistance band standing internal rotation at 90 degrees <p>Motor Control</p> <ul style="list-style-type: none"> ▪ 90/90 ball dribbles, over-head soccer throws, medicine ball chest pass, prone ball drops, standing ball drops, 90/90 over the shoulder eccentric catch and throw, body blade
Criteria to Progress	<ul style="list-style-type: none"> ▪ No pain or tenderness ▪ 5/5 shoulder strength ▪ Satisfactory shoulder stability ▪ Use Quick DASH and/or PENN shoulder scale ▪ Upper Extremity Functional Assessment <ul style="list-style-type: none"> ○ Full pain-free PROM and AROM ○ Joint position sense < 5-degree margin of error ○ Strength 85% of uninvolved arm with isokinetic testing or handheld dynamometer ○ ER/IR ratio > 64% ○ Scapular dyskinesis test symmetrical ○ Functional performance and shoulder endurance tests > 85% of uninvolved arm ○ Males > 21 taps; females > 23 taps on CKQUEST ▪ Additional UE Functional Tests <ul style="list-style-type: none"> ○ One-arm hop test ○ Push-up test ○ BABER

PHASE VII: UNRESTRICTED RETURN-TO-SPORT (6+ MONTHS AFTER SURGERY)

Rehab Goals	<ul style="list-style-type: none"> ▪ Maintain pain-free ROM ▪ Gradual return to strenuous work activities ▪ Gradual return to recreational activities ▪ Gradual return to sports activities
Additional Interventions *Continue with Phase II-VI interventions	<p>Strengthening</p> <ul style="list-style-type: none"> ▪ 50% 1 RM bench press, progress slowly (coordinate with physician) ▪ See specific return-to-sport program (coordinate with physician)

Evan D. Boyd, MD
Knee, Shoulder, & Sports Medicine Surgeon
518-489-2663 ext 5215

Return-to-Sport	<ul style="list-style-type: none">▪ For the recreational or competitive athlete, return-to-sport decision making should be individualized and based upon factors including level of demand on the upper extremity, contact vs non-contact sport, frequency of participation, etc. We encourage close discussion with the referring surgeon prior to advancing to a return-to-sport rehabilitation program.
------------------------	--

**Acknowledgement: This rehab protocol was largely adopted from the protocols at MGH Sports Medicine Physical Therapy, which can be found at <https://www.massgeneral.org/orthopaedics/sports-medicine/physical-therapy/sports-rehab-protocols>*